

Heterozygous OTX2 deletion in a boy with normal eye development and normal pituitary function

Boros E¹, Boitsios G², Vilain C^{3,4,5}, Balikova I⁶, Heinrichs C¹, Brachet C¹

¹ Paediatric Endocrinology Unit, Hôpital Universitaire des Enfants Reine Fabiola, Université Libre de Bruxelles, Brussels, Belgium; ² Pediatric Radiology Unit, Hôpital Universitaire des Enfants Reine Fabiola, Université Libre de Bruxelles, Brussels, Belgium; ³ Department of Genetics, Hôpital Universitaire des Enfants Reine Fabiola, ULB Center of Human Genetics, Université Libre de Bruxelles, Brussels, Belgium; ⁴ Department of Genetics, Hôpital Erasme, ULB Center of Human Genetics, Université Libre de Bruxelles, Brussels, Belgium; ⁵ Interuniversity Institute of Bioinformatics in Brussels, Université Libre de Bruxelles, Brussels, Belgium; ⁶ Pediatric Ophthalmology Unit, Hôpital Universitaire des Enfants Reine Fabiola, Université Libre de Bruxelles, Brussels, Belgium

Introduction. *Orthodenticle homeobox 2 (OTX2)* is a transcription factor that plays a critical role in forebrain and eye development. It likely interacts with HESX1 in Rathke's pouch and regulates various transcription factors as RX1, PAX6, LHX2. OTX2 heterozygous mutations or deletions lead to eye malformations such as anophthalmia, microphthalmia, coloboma, optic nerve hypoplasia but also dystrophy of the retinal pigmented epithelium (role in the mature retina) (Henderson and al), normal or hypoplastic pituitary gland and normal or ectopic posterior pituitary gland with isolated growth hormone deficiency or combined pituitary hormone deficiency (Diaczok and al). There is no genotype-phenotype correlation and variable presentation in the same family with whole gene deletion has already been described (Jones E and al).

Case report:

A 3,2 year-old boy presented to our pediatric endocrinology unit for short stature.

Neonatal history:

- born at term after an uneventful pregnancy with 3,010 kg and 50 cm

Medical history:

- familial Mediterranean fever

Familial history:

The mother's height is 156,3 cm and the father's height is 162 cm (target height 165 cm +/-8,5 cm (-1,7 SDS), Parents are non-consanguineous and healthy, Caucasians



First examination

Height was 86,4 cm (-2,8 DS),

Weight 11,05 kg


Head circumference 46,8 cm (-3,4 DS).

No dysmorphic features. Tanner stage A1P1G1 with 2 ml testis bilaterally



Laboratory analysis:

- Normal thyroid function: free T4 18,6 pmol/l (reference range: 12,3-22,8)
- IGF1 normal for age : 67 ng/ml and 71 ng/ml (reference range: 15-200)
- insulin tolerance test showed a normal response for both growth hormone and cortisol : GH peak 7,44 ng/ml (normal value >7) and cortisol peak 561 nmol/l (normal value >550).
- Microarray analysis showed a heterozygous deletion of 14q22.3 including the whole OTX2 gene
- Both parents had normal microarray results



Cerebral MRI shows a malformation of the pituitary region, with an almost absent sella turcica, normal size but rounded anterior pituitary, with an ectopic posterior pituitary gland, thin pituitary stalk and no optic nerve abnormality.

Extensive ophthalmological examination showed no eye malformation or vision problem in our patient.

In conclusion we present on the case of a boy with heterozygous OTX2 deletion who harbors flat sella turcica, ectopic posterior pituitary but neither hormonal deficiencies so far nor eye malformation.

The anterior pituitary function will need to be followed-up as evolving hormonal deficiency is possible.

The retinal function will also need to be followed-up given the reported role of OTX2 not only in the developing, but also in the mature retina.

References:

- Henderson RH, Williamson KA, Kennedy JS, Webster AR, Holder GE, Robson AG, FitzPatrick DR, van Heyningen V, Moore AT. A rare de novo nonsense mutation in OTX2 causes early onset retinal dystrophy and pituitary dysfunction. Mol Vis. 2009 Nov 21;15:2442-7; Jones GE, Robertson L, Warman P, Craft EV, Cresswell L, Vasudevan PC. 14q22.3 Microdeletion encompassing OTX2 in a five-generation family with microphthalmia, pituitary abnormalities, and intellectual disability. Ophthalmic Genet. 2016 Sep;37(3):352-3; Diaczok D, Romero C, Zunich J, Marshall I, Radovick S. A novel dominant negative mutation of OTX2 associated with combined pituitary hormone deficiency. J Clin Endocrinol Metab. 2008 Nov;93(11):4351-9; Beby F, Lamonerie T. The homeobox gene Otx2 in development and disease. Exp Eye Res. 2013 Jun;111:9-16.; Ragge NK, Brown AG, Poloschek CM, Lorenz B, Henderson RA, Clarke MP, Russell-Eggitt I, Fielder A, Gerrelli D, Martinez-Barbera JP et al. Heterozygous mutations of OTX2 cause severe ocular malformations. Am J Hum Genet. 2005 Jun;76(6):1008-22.

