

Serum Fibroblast Growth Factor 23 And Klotho Concentrations In Children And Adolescents With Obesity

S. KARAMPATSOU¹, S. GENITSARIDI¹, P. KASSARI¹, G. KOURLABA², E. CHARMANDARI^{1,3}

1. Division of Endocrinology, Metabolism and Diabetes, First Department of Pediatrics, National and Kapodistrian University of Athens Medical School, 'Aghia Sophia' Children's Hospital, Athens, Greece

2. Center for Clinical Epidemiology and Outcomes Research (CLEO), Athens, Greece

3. Division of Endocrinology and Metabolism, Center of Clinical, Experimental Surgery and Translational Research, Biomedical Research Foundation of the Academy of Athens, Athens, Greece



www.childhood-obesity.gr

INTRODUCTION

Obesity in childhood and adolescence represents one of the main health problems of the 21st century. Fibroblast Growth Factor 23 (FGF-23) and its co-receptor, Klotho, play an important role in mineral metabolism. However, little is known about their role in obesity.

AIM


Our aim was to determine the concentrations of FGF-23 and Klotho in children and adolescents with overweight and obesity.

METHOD


Three hundred and twenty subjects [163 males (50.9%), 157 females (49.1%), 158 prepubertal (49.4%), 162 pubertal (50.6%)], aged [mean \pm standard deviation (SD)] 10.5 \pm 2.8 years, were studied prospectively for one year. Subjects were classified as obese (n=211, 66%), overweight (n=99, 30.9%) and of normal BMI (n=10, 3.1%) and were enrolled in a one-year lifestyle intervention program of diet and physical exercise. Body composition and biochemical and endocrinologic parameters were determined at the beginning and at the end of the study. The study was approved by the Committee on the Ethics of Human Research, and written informed consent was obtained

RESULTS

- **Significant decrease in BMI ($p=0.001$), total cholesterol ($p=0.001$), LDL ($p=0.001$), HBA1C ($p=0.021$) and FGF-23 ($p=0.004$) concentrations, and percentage of body fat ($p=0.001$).**
- **Significant increase in HDL ($p=0.001$), parathormone (PTH) ($p=0.001$), 25-OH-Vitamin D (VitD) ($p=0.003$), adiponectin ($p=0.001$) and Klotho ($p=0.001$) concentrations, and percentage of muscle mass (PMM) ($p=0.001$).**
- **Klotho concentrations correlated negatively with the waist-to-hip ratio ($p=0.001$) and VitD ($p=0.022$), and positively with PTH ($p=0.050$).**
- **The change in FGF-23 concentrations correlated positively with the change of Apolipoprotein1 (Apo1) ($r=0.200$, $p=0.001$).**
- **The change in Klotho concentrations correlated positively with the change in BMI ($r=0.169$, $p=0.009$), diastolic blood pressure ($r=0.267$, $p=0.022$), waist circumference ($r=0.319$, $p=0.001$), Apo1 ($r=0.171$, $p=0.011$), HOMA-IR ($r=0.259$, $p=0.001$), PMM ($r=0.226$, $p=0.004$) and adiponectin ($r=0.215$, $p=0.002$).**
- **The change in BMI correlated positively with Klotho change in males ($r=0.438$, $p=0.001$).**
- **The change in BMI correlated positively with the change in FGF-23 ($r=0.364$, $p=0.023$) and Klotho ($r=0.462$, $p=0.001$) concentrations in subjects that entered puberty.**
- **Participants had normal kidney function.**



The change of FGF-23 and Apolipoprotein1 before and 12 months after the intervention



The change of Klotho and HOMA-IR before and 12 months after the intervention

	BMI change	FGF-23 change r / rho (p-value)	Klotho change r / rho (p-value)
Gender			
Males		0,034 (0,683)	0,438 (0,001)
Females		-0,028 (0,738)	-0,046 (0,625)
Pubertal stage (after intervention)			
Same pubertal status (prepubertal or pubertal)		-0,074 (0,289)	0,127 (0,099)
Subjects that entered puberty		0,364 (0,023)	0,462 (0,001)

CONCLUSIONS

These findings indicate an association between bone-derived FGF-23, its co-receptor Klotho, obesity and cardiovascular risk factors. More studies are needed in order to determine the role of FGF-23 and Klotho as biomarkers of childhood obesity, especially in adolescence.

REFERENCES

- 1 Zaheer S et al. Fibroblast Growth Factor 23, Mineral Metabolism, and Adiposity in Normal Kidney Function. J Clin Endocrinol Metab. 2017; 102(4); 1387-1395
- 2 Xu Y et al. Correlations between serum concentration of three bone-derived factors and obesity and visceral fat accumulation in a cohort of middle aged men and women. Cardiovasc Diabetol. 2018; 17(1); 1-9
- 3 Gkentzi D et al. Fibroblast growth factor 23 and Klotho serum levels in healthy children. Bone. 2014; 66(2014); 8-14

ACKNOWLEDGEMENTS

We would like to express our appreciation to both the participants and the staff of the Out-Patient Clinic for the Prevention and Management of Overweight and Obesity.

CONTACT INFORMATION

Email: karampatsousi@gmail.com

Telephone number: +306972861400

