

FRUCTOSAMINE LEVEL IN TYPE 1 DIABETES MELLITUS CHILDREN PERFORMING RAMADHAN FASTING



Muhammad Faizi, Nur Rochmah

Department of Child Health Faculty of Medicine Airlangga University-Dr. Soetomo General Hospital Surabaya

Abstract:

Background: Ramadan fasting may influence metabolic control in Type 1 Diabetes Mellitus. Fructosamine is an accurate metabolic control in a short-term period. Comparison of fructosamine between intensive and conventional insulin regimen in T1DM children has not been widely studied.

Methods:

- Observational analytic study
- conducted in endocrine-OPC Dr.Soetomo hospital
- during Ramadan 2013 and 2015.
- Inclusion criterias: T1DM children performed Ramadan fasting before, and already diagnosed >6 months before the study.
- **Exclusion criterias:**
- severe hypoglycemia<3 months before the study,
- recurrent hypoglycemia(blood glucose <65mg/dl)
- hyperglycemia with DKA<3months before study conducted
- hospitalized patients
- History of kidney and liver disease and severe malnutrition.
- Fructosamine was evaluated 3 times (one week before, in the middle and at the end of Ramadan fasting).
- Daily blood glucose monitoring, DKA events, and hypoglycemia episodes were observed.
- Patients were divided into 2 groups: group 1 given intensive-treatment (Detemir, shortacting insulin) and group 2 conventional (intermediate, short-acting insulin).

Results:

Table 1. Demographics Data Children with T1DM

Characteristics	Frequency n (%)	Mean	Range	SD
Gender				
Male	10 (41.7)			
Female	14 (58.3)			
Age (year old)		14.92	9-19	2.6
5 – 9 years old	1 (4.2)			
10 – 14 years old	8 (33.3)			
15 – 19 years old	15 (62.5)			
Weight (kg)	` '			
Before Ramadhan		45.3	27-63	9.5
End of Ramadhan		44.6	26-63	9.4
fasting				
Length of Suffering From		4.5	0.5 - 9	2.5
T1DM (year)				
History of DKA		2.13	1-3	
1 time	3 (12.5)			
2 times	15 (62.5)			
3 times	6 (25)			
Insulin Regimens				
Intensive	9 (37.5)			
Conventional	15 (62.5)			

Objective:

To compare fructosamine level between intensive and conventional insulin regimens during Ramadan Fasting in T1DM.

Table 2. The Comparison of Fructosamine Levels Before, Middle, and at The End of Ramadhan Fasting

Periode of Ramadhan Fasting	Mean of F	Mean of Fructosamine level (µmol/L)				
	Intensive	Conventional	р			
Before	497.1 (± 146.6)	531,2(± 126.9)	0.5*			
Middle	458.2 (± 104.8)	529,6(± 118.6)	0.1*			
End	467 (± 118)	512(± 141.5)	0.4*			

Comparison of Fructosamine level between groups before (497.1vs.531.2 µmol/L,P=0.5), middle (458.2vs.529.6 µmol/L,P=0.1) and at the end of Ramadan(467vs.512 µmol/L,P=0.4) were not significant(P>0.05).

Most subjects (91.6%) had hyperglycemia without DKA with a mean of frequency 17.3 times.

Neither symptomatic hypoglycemia nor DKA was found in this study.

Conclusion:

Fructosamine level during Ramadan fasting between intensive and conventional insulin regiments were similar. Type-1 DM fasting children did not experience symptomatic hypoglycemia or DKA.

Keywords:

fructosamine, metabolic control, ramadan fasting, insulinfructosamine, metabolic control, T1DM, ramadan fasting, insulin

REFERENCES

1. El-Hawary A, Salem N, Elsharkawy A, et al. Safety and metabolic impact of Ramadan fasting in children and adolescents with type 1 diabetes. J Pediatr Endocrinol and Metabolism. Walter de Gruyter 2016;29(5).

2. AlAlwan I and Banyan AA. Effects of Ramadan fasting on children with Type 1 diabetes. International Journal of Diabetes Mellitus 2010;2(2):127–129.

3. Nansseu JRN, Fokom-Domgue J, Noubiap JJN, et al. Fructosamine measurement for diabetes mellitus diagnosis and monitoring: a systematic review and meta-analysis protocol. BMJ Open 2015;5(5):7689–7689.

Poster Number: 113-P3

The Topic: Diabetes and Insulin, 1st Author: Muhammad Faizi











